

# **A Research of the Relationship between Childhood Maltreatment and Suicide: Does Support Increase The Risk of Suicide in Maltreated Children?**

C. Gilde (3269310)

2011 - July

## **Abstract**

This research focuses on the relationship between childhood maltreatment and suicide. The types of maltreatment that were included are physical abuse, sexual abuse, emotional maltreatment and physical neglect, and suicide was operationalized by the factors suicidal thoughts, suicide attempts and a combination of both. It was hypothesized that the types of maltreatment were all related to the suicide variables, and that these relationships are influenced by the variables negative self-concept, social support, teacher support, family support, relationship with parents, and present contact with parents. Results showed that there was hardly any significant direct relationship between the maltreatment types and suicide variables. Only sexual abuse was related with suicide attempts and suicide (total), and emotional maltreatment with suicide (total). When several risk/protective factors were added in a logistic regression analysis, 'negative self-concept' and teacher support were the most important risk factors, and 'present contact with parents' was the biggest protective factor of suicide in maltreated persons.

Utrecht University, Netherlands  
Master Clinical and Health Psychology  
Supervisor: Prof. Dr. R. Kleber

Videnscenter for Psykotraumatologi  
Odense, Denmark  
Supervisor: Prof. A. Elklit

## **Foreword**

For the master Clinical and Health Psychology at Utrecht University, I wrote this master thesis the first semester of my master. A couple of months before the master started, I decided to try to do my master thesis abroad. Together with Mireia van der Vegt, I sent motivational- and application letters to institutes that are focused on trauma. Prof. Ask Elklit from the Videnscenter for Psykotraumatologi in Odense wrote that we were welcome to come over and do our thesis at the centre. There was a large database with some interesting data. Because of our interest in trauma, we choose the subject of childhood maltreatment in combination with suicide.

In the first place, I would like to thank Mireia van der Vegt for our great cooperation. I also want to thank my both supervisors. I thank Prof. Dr. Rolf Kleber from Utrecht University and Prof. Ask Elklit from Syddansk Universitet for their professional supervision and advises, and for making it possible for me to do my thesis abroad. Furthermore, I would also like to thank Prof. Ask Elklit and the Videnscenter for Psykotraumatologi in Odense for their warm welcome and hospitality. Additionally, I am grateful to The Danish National Centre for Social Research for collecting all data, and for letting us use the database. Finally, I would like to thank everybody who helped and supported me in the process of my master thesis.

## Index

Abstract.....	1
Foreword.....	2
Introduction.....	4
Methods.....	9
Results.....	15
Discussion.....	21
References.....	26
Appendix.....	30

## **Introduction**

### *Child maltreatment*

Child maltreatment is in the Western world a very common form of violence perpetrated by caregivers (Hahm, Lee, Ozonoff, & Van Wert, 2010). It is an important psychological issue that needs attention in order to minimize the negative consequences. According to the World Health Organization, 25-50% of globally all children report being physically abused. Furthermore, approximately 20% of women and 5-10% of men report being sexually abused as children. However, the so far reported prevalence rates of child maltreatment often differ. The WHO defines that child maltreatment, sometimes referred to as child abuse and neglect, includes all forms of physical and emotional ill-treatment, sexual abuse, neglect and exploitation that results in actual or potential harm to the child's health, development or dignity. When abuse and neglect occur to children under 18 years of age, it's called child maltreatment (WHO, 2011). In this study, the focus will be on the relationship between the specific types of child maltreatment and suicide, and factors that influence this relationship.

### *Consequences child maltreatment*

Many negative outcomes have been found for child maltreatment in research studies. Child maltreatment causes one of the most adverse and stressful challenges that confronts children (Cicchetti, Rogosch, Sturge-Apple, & Toth, 2010). Literature has shown a relationship between being a victim of sexual abuse, neglect, physical, or emotional maltreatment, and post-traumatic stress disorder (PTSD) (Burns, Jackson, & Harding, 2010; Cantón-Cortés & Cantón, 2010). For this disorder, women show a higher prevalence rate than men (Ditlevsen & Elklit, 2010). PTSD consists of many symptoms, including anxiety, disturbances of memory, elevated arousal, avoidance and fear of horror. These symptoms persist for prolonged periods after an unusually distressing experience (Rachman, 2004). Scott, Smith, and Ellis (2010) found significant associations between prospectively ascertained child maltreatment and a number of mental disorders. The associations were strongest for post-traumatic stress disorder and obsessive-compulsive disorder. High levels

of trauma-related symptoms were also consistently associated with a history of childhood sexual abuse (Aspelmeier, Elliott, & Smith, 2006).

Additionally, according to Burns et al. (2010), there is preliminary support for greater emotion regulation difficulties among emotionally abused women. They found that the relationship between childhood maltreatment (physical or emotional) and post-traumatic symptoms was partially explained by emotion regulation difficulties. This study indicated that emotional abuse is a stronger predictor of emotion dysregulation, when compared to sexual or physical abuse. Kim and Cicchetti (2010) found that more difficulties in emotion-regulation and greater externalizing problems were reported by maltreated children – especially for the subtypes of neglect, physical, and/or sexual abuse, multiple subtypes, and earlier onset. For emotional maltreatment, the relation with emotion dysregulation was not significant, but also not negligible.

Several other negative outcomes of child maltreatment are found. Hahn et al. (2010) have demonstrated that women making the transition from adolescence to young adulthood, are affected negatively and sustainably on many developmental outcomes by child maltreatment. They found that women who were maltreated in several ways are – when compared to women who experienced one type of maltreatment – more likely to report running away from home and to report Sexually Transmitted Disease diagnosis. Furthermore they were more likely to have had sex for money, and to have had sex before the age of 15.

Specific results have been found for the consequences of sexual abuse in some studies. Senn and Carey (2010) showed that childhood sexual abuse is uniquely associated with adult sexual risk behavior. Lower levels of attachment security in close-adult, parent-child and peer relations, were associated with a history of childhood sexual abuse (Aspelmeier et al., 2007).

In addition to all of these consequences, child maltreatment has often been linked to suicide, making it of special interest for this study. A theory that could explain the death wish in maltreated persons is the interpersonal-psychological theory. According to this theory, the two interpersonal states ‘perceived burdensomeness’ and ‘thwarted belongingness’

confluence. Those states represent feeling like one's existence is a burden of friends, family members and/or society, and feeling alienated from friends, family, or other valued social circles. The states are independently related to elevated risk for the desire for suicide, but if they co-occur the risk is greatest (Ribeiro & Joiner, 2009). In this research, it is assumed that childhood maltreatment is related to these two states, and therefore that maltreatment is related to suicide. One of the research questions of the current research will be: Does childhood maltreatment increase the risk of suicidal thoughts and suicide attempts?

### *Child maltreatment & suicide*

According to Meadows and Kaslow (2002), the increased risk for many psychological difficulties resulting from child maltreatment, may precede the development of suicidal behaviors. In a research of Bruffaerts et al. (2010), in 21 Western (including the Netherlands) and non-Western countries, it was revealed that physical and sexual abuse were strongest related with suicidal risk behavior. However, this research did not include emotional maltreatment as a type of maltreatment. Brown, Cohen, Johnson, and Smailes (1999) and Dube, Anda, Felitti, Chapman, Williamson, and Giles (2001) found that adverse childhood experiences increased the risk of ever attempting suicide from 2- to 5 fold. The risk of ever attempting suicide and of attempted suicide increased dramatically according to the latter research, when the number of such experiences increased. Their analysis suggested that the abusive or traumatic childhood experiences attribute to approximately two thirds (67%) of suicide attempts. Another study found that a history of childhood maltreatment is likely to play a key role in the onset and recurrence of suicide attempts (Perroud et al., 2007). For the severity of suicidal behavior, only sexual abuse played a key role. Similar findings are reported by studies focusing solely on women by Hahm et al. (2010) and Sfoggia, Pacheco, and Grassi-Oliveira (2008). According to Hahm et al., women who were maltreated in several ways are – when compared to women who experienced one type of maltreatment – more likely to report suicide ideation and attempted suicide.

As mentioned above, a history of maltreatment has a significant association with suicidal ideation. Suicidal ideation seems to be already evident in children as young as 8

years old. Maltreated children approximately have a 2-fold risk to report suicidal ideation at this age. This increased risk is associated with the severity of physical abuse and with the presence of multiple types of maltreatment (Thompson et al., 2005). Additionally, Cheng et al. (2009) found that suicidal ideation and attempts increase with age and Brown et al. (1999) indicate that adolescence seems to be the most vulnerable period for suicide in sexually abused youths.

Looking at the influence of specific types of maltreatment on the relation with suicide, Ryan, Kilmer, Cauce, Watanabe and Hoyt (2000) found minor differences between three subtypes of maltreatment. They compared the types physical abuse only, sexual abuse only and both types of abuse, with no reported abuse. The combination of physical and sexual abuse showed highest risk of attempted suicide. Moreover, physical abuse only and sexual abuse only gave a higher risk of attempted suicide when compared to the 'no reported abuse' group. In another study, the relationship between different kinds of maltreatment has been investigated as well. The research showed higher levels of suicide proneness for individuals that were neglected in their childhood, than those who reported physical abuse alone or sexual abuse alone (Arata, Langhinrichsen-Rohling, Bowers, & O' Brien, 2007). In the research of Brown et al. (1999) results showed that childhood neglect alone was presumably not responsible for suicidal behavior, since its effects couldn't be separated from those of other risk factors. Here, sexual abuse involved the greatest risk of suicide, independently of the contextual risks under which the abuse occurs. In the mentioned study, physical abuse alone contributed to repeated suicide attempts during adolescence as well.

In the current research, the relation between the four types of maltreatment and suicidal thoughts and attempts will be examined. Because of the differences in sensitivity for contextual risks, this study will also focus on factors affecting the association between childhood maltreatment and suicide.

## *Factors affecting the association*

### Risk factors

Research has shown that several factors affect the relationship. First, the risk factors will be discussed. In a sample of low-income African American women, Meadows and Kaslow (2002) found that hopelessness in adulthood partially mediates the relation between a childhood maltreatment history and adult suicide attempts. This could be explained by the suggestion of Schotte and Clum (1982) that interpersonal poor problem solving in conjunction with negative life stress might lead to feelings of hopelessness, which in turn result in the development of suicide ideation and intent. This in accordance with the previously mentioned interpersonal-psychological theory (Ribeiro & Joiner, 2009). Additionally, Salzinger, Rosario, Feldman, and Ng-Mak (2007) discovered that combined childhood and adolescent internalizing problems (depression) mediate the effect of preadolescent abuse on suicidal attempts. Furthermore, they found that continued parental abusive behavior (physical) in adolescence is significantly associated with suicidal behavior. In this sample, internalizing problems in adolescence contributed to increased risk of suicidal behavior. Conversely, stressful life events did not have any significant influence on the risk of suicidal behavior. However, parental divorce plays a role in the association according to Boxer (2010). He has found an additive effect between parental divorce and child abuse for suicide attempts. Notably, further adjusting it for parental psychopathology caused the effect to be reduced.

Moreover, Locke and Newcombe (2005) found in a sample of Latino males that several hypothesized domains did not predict suicidality: physical abuse, physical neglect, emotional neglect, father's alcohol-related problems, childhood quality, good relationships with family, assertiveness, competence, religious commitment, cigarette use, alcohol use and marijuana use.

### Protective factors

Some protective factors have also been found for suicidality in previous research. Attachment security protects at least partially against negative outcomes of childhood sexual abuse (Aspelmeier et al., 2007). Infants who have secure relationships with their caregivers use them as a source of comfort and support to control feelings of distress when they arise. For these infants, the caregiver's presence signals safety and is an unequivocal source of comfort (Simpson & Rholes, 1998). According to Salzinger et al. (2007), parental attachment acts as a protective factor for the risk of suicidal behavior, whereas attachment to friends doesn't have any significant influence on the risk of suicidal behavior. Locke and Newcomb (2005) found that a good relationship with parents, problem-solving confidence and being law abiding act as protective factors. Moreover, Eisenberg, Ackard, and Resnick (2007) concluded that family connectedness has a strong protective association with suicidal

ideation and attempts. Other significant protective factors for both male and female adolescents were teacher caring, other adult caring and school safety. In conclusion, family seems to play an important protective role. Therefore, family factors (e.g. family support, relationship with parents, contact with parents) will be closely examined in this research. Contact with parents has not yet been included in research on the topic. Because this factor is about specific behavior that doesn't concern feelings, it is included in this research, which makes the study unique.

### *Suicide*

There are other factors that affect the risk of suicidal behavior, which are not yet related to childhood maltreatment. In order to conduct a complete as possible view on the subject, these factors should be considered as well. According to the interpersonal-psychological theory, the misperception that the self is so incompetent that one's existence is a burden on family members, friends, or other valued social circles, leads to the increased risk of the belief that one's death is worth more than one's life (Ribeiro & Joiner, 2009). A research of Kaslow et al. (2002), which focused on low-income abused African American women, shows that being hopeful, being high in self-efficacy, having the capacity to obtain resources, having adaptive coping skills and strong social support significantly reduce the risk for attempting suicide in this sample. In a Turkish study, self-esteem was a significant predictor of suicidal risk scores. Greater suicidal ideation and suicide potential showed a significant correlation with lower self-rated problem-solving skills (Eskin, Ertekin, Dereboy, & Demirkiran, 2007). These studies lead to the inclusion of the factor negative self-concept in the current research on the relation between childhood maltreatment and suicide. On the subject of the influence of gender on suicide, literature is not consistent. Eskin et al. (2007) found that females had a higher risk for suicide ideation, but not for suicide attempts. In contrast, Dube et al. (2001) and Fleming Merry, Robinson, Denny and Watson (2007) showed that women had an increased risk for suicide attempts. Moreover, no gender differences were found for suicidal risk behavior when maltreatment was also taken into account (Dube et al., 2001, Thompson et al., 2005). Because of these inconsistencies, gender will also be included in this research. Furthermore, family support seems to play an important protective role in suicidal risk behavior (Au, Lau, & Lee, 2009; Randell, Wang, Herting, & Eggert, 2006; Sharaf, Thompson, & Walsh, 2009). According to Compton, Thompson, and Kaslow (2005), lower reported levels of family adaptability, family cohesion, social embeddedness and social support increased the relative rate of suicide attempt. In another study, depressive symptoms, alcohol abuse, non-heterosexual orientation and exposure to suicide behavior by others were associated with an increased risk of suicide attempts (Fleming et al., 2007). This study also found that a caring home and a fair and safe school environment contributed to a lower risk for suicide attempts. Additionally, Dervic,

Oquendo, Grunebaum, Ellis, Burke, and Mann (2004) showed in their research that religion was inversely related to suicidal ideation, though the effect was mediated by a greater moral objection to suicide which may have represented traditional religious beliefs. Thus, many factors influence the risk of suicide.

#### *Aim of this study*

This study will investigate whether childhood maltreatment alone is responsible for suicidal behavior, or whether the association between childhood maltreatment and suicidal thoughts and attempts is influenced by internal/external factors. The influence of negative self-concept, support (social, teacher, family), relationship with parents, and present contact with parents will be investigated.

This research is unique in the fact that it makes use of a large sample, partially consisting of child protection cases. The implementation of some factors that have not yet been proven to influence the association between childhood maltreatment and suicide, also makes this study exclusive. The research questions are: *Does childhood maltreatment influence the risk of suicidal thoughts and suicide attempts, and is there a difference for the several types of maltreatment? Which factors influence the relation between childhood maltreatment and suicide?* First, the relationships between the types of childhood maltreatment and suicide, suicidal thoughts and suicide attempts will be researched. It is hypothesized that experiencing any of the types of childhood maltreatment would be associated with a higher risk of reporting suicide, suicidal thoughts and suicide attempts. Furthermore, the relations between all types of maltreatment and the factors 'negative self-concept', 'social support', 'teacher support', 'family support', 'relationship with parents' and 'present contact with parents' will be investigated. Similarly, the relation between the suicide variables and the factors will be examined. Finally, the effects of the factors for the relationship between maltreatment and suicide will be studied. It is expected that a 'negative self-concept' increases the risk of suicidal behaviors in all types of maltreatment. On the contrary, it is assumed that 'social support', 'family support', 'teacher support', 'relationship with parents', and 'present contact with parents' act as protective factors of suicidal behaviors.

## **Methods**

### *Design*

The design used in this research is a between-subjects design. Questionnaires were used (see Appendix).

### *Participants*

A total of 2980 interviews were achieved with a response rate of 63%. The most common reasons for not participating were refusal to take part in the study (21%), lack of

contact (13%), and illness or disability (2%). To adjust for the oversampling of child protection cases the data have been weighted so that findings are representative of the total Danish population of young people aged 24 years.

The demographic characteristics of the total sample (weighted), the child protection cases, and the non child protection cases are summarized in Table 1. Fifty-two percent of the total sample (weighted) was male.

Table 1 *Demographic Characteristics of Non Child Protection Cases and Child Protection Cases.*

	Total Sample weighted N = 2980 (%)	Non child protection case N = 2128 (%)	Child protection case N = 852 (%)
<i>Gender</i>			
• Male	1555 (52.2)	1106 (52.0)	473 (55.5)
• Female	1425 (47.8)	1022 (48.0)	379 (44.5)
<i>Living Accommodation</i>			
• Home owner	837 (28.1)	608 (28.6)	176 (20.7)
• Rented accommodation	1949 (65.6)	1387 (65.3)	587 (69.3)
• Supported living	175 (5.9)	123 (5.8)	62 (7.3)
• Other	11 (0.4)	5 (0.2)	22 (2.6)
<i>No. of Dependants &gt;18</i>			
• 0	2717 (91.3)	1954 (91.9)	695 (81.7)
• 1	197 (6.6)	132 (6.2)	110 (12.9)
• 2	52 (1.8)	33 (1.6)	41 (4.8)
• =>3	10 (0.3)	7 (0.3)	5 (0.6)
<i>Education</i>			
• No Higher education	507 (17.1)	324 (15.3)	372 (44.0)
• Higher/ vocational education student	1390 (46.9)	1024 (48.4)	209 (24.7)
• Completed higher/vocational education	1066 (36.0)	768 (36.3)	264 (31.2)
<i>Employment</i>			
• Employed	1611 (54.1)	1141 (53.6)	519 (61.2)
• Unemployed	152 (5.1)	87 (4.1)	173 (20.4)
• Student	1190 (40.0)	882 (41.5)	147 (17.3)
• Other	24 (0.8)	17 (0.8)	9 (1.1)
<i>Marital Status</i>			
• Married	184 (6.2)	128 (6.0)	74 (8.7)
• Cohabiting	1186 (39.8)	852 (40.0)	307 (36.1)
• Single	1610 (54.0)	1148 (53.9)	470 (55.2)

Note. All values are weighted.

### *Procedure*

Data were collected by The Danish National Centre for Social Research in 2008 and 2009. The study used a stratified random probability sample of young people aged 24 years. The study was funded by the Danish Research Council. A sample of 4718 young adults with Danish citizenship was randomly selected by Statistics Denmark using the total birth cohort of all children born in 1984 (excluding persons who had refused to participate in national research or were imprisoned). Participation in the study was entirely voluntary and the study was approved by the Danish Data Protection Agency. To increase the number of participants who had experienced childhood abuse and neglect, children who had been in child protection were oversampled by stratifying the number of “child protection cases”

versus “no child protection cases” (1/3:2/3). A child protection case was defined as a case where the council (according to the files of local social workers) had provided support for the child and the family or placement with a foster family due to concerns about the well-being and development of the child.

The data were collected using a structured interview, which was conducted as a telephone interview or as a residential interview according to the participants' choice. The average duration of the interview was estimated to 43 minutes. The response format was precoded, but with an option for respondents to add additional comments if necessary. The interview did not define abuse and neglect but asked respondents if they had experienced specific behaviors towards them. A letter sent prior to the data collection informed each participant about the nature of the research, the possibility of being interviewed at home, and the procedures securing confidentiality. Persons who did not respond to the letter, were contacted by telephone if possible, and then were contacted at their home address.

The interviewers were carefully trained by The Danish National Centre for Social Research prior to the data collection. The training included detailed oral information and standardized written instructions regarding the purpose and content of the study. Moreover, test trials were conducted to familiarize the interviewers with the questionnaire and the coding procedure. The study included several sensitive questions regarding sexual abuse and violence. Hence, participants who were interviewed in their home, answered these questions using computer-assisted personal interviewing, whereby respondents could enter their answers directly on to a laptop computer. This method has been validated in similar studies (e.g. May-Chahal & Cawson, 2005). Furthermore, all participants were offered a telephone number of an experienced psychologist at the end of the interview, in case they wanted any counseling help.

### *Measures*

Physical Abuse. Retrospective reports on physical abuse from parents or guardians were obtained utilizing single-items that describe different experiences of serious violent treatment (see Table 2). The items asked whether respondents had experienced seven different types of abuse and scores were rated on a two-point (yes/no) format.

Table 2 *Experience of physical abuse from parents / guardians by sex and child protection status*

	Total	Male	Female	Non Child protection case	Child protection case
Total (N)	2980	1579	1401	2128	852
Weighted (N)	2980	1555	1425	2794	186
Beaten with an object, such as a whip or coat hanger? (%)	2.7	3.2	2.1	2.1	11.5
Threatened with a weapon, such as a knife or a gun? (%)	0.4	0.4	0.5	0.3	2.5
Had objects thrown at you? (%)	2.7	3.2	2.2	2.1	12.1
Grabbed round the neck and choked? (%)	0.5	0.3	0.6	0.3	2.5
Been left with burn or bite marks? (%)	0.2	0.1	0.3	0.1	1.9
Had injuries such as broken bones, stab wounds, brain haemorrhage, or burns which were treated by a doctor? (%)	0.3	0.2	0.4	0.2	2.3
Been hit, kicked or exposed to violence which has resulted in bruising, bleeding, or other physical injuries? (%)	1.6	1.6	1.6	1.1	9.6
Total (%)	5.2	6.1	4.1	4.2	19.8

*Note.* All percentage values are weighted; Total = Positive endorsement of at least one item; Categories were not mutually exclusive.

Sexual Abuse. Retrospective reports on sexual abuse from parents or guardians were obtained utilizing single-items that describe experiences of serious sexual abuse under the age of 13 (see Table 3). Only cases who confirmed that they had been exposed to sexual abuse were asked about the four types of abuse, and scores were rated on a two-point (yes/no) format.

Table 3 *Experiences of sexual behavior under the age of 13*

	Total	Male	Female	Non Child protection case	Child protection case
Total (N)	2980	1579	1401	2128	852
Weighted (N)	2980	1555	1425	2794	186
Childhood Sexual Abuse (Y/N)	1.0	0.3	1.8	0.7	6.3
Experienced sexual touching or someone exposing their private parts /sex organs to you (%)	2.6	0.5	4.8	2.1	9.5
Experienced attempted intercourse (%)	2.6	0.4	5.0	2.2	9.2
Experienced forced / completed intercourse (%)	1.9	0.2	3.8	1.6	6.5
Experienced other types of sexual behaviour (%)	0.8	0.2	1.4	0.6	3.8

*Note.* All percentage values are weighted; Only those cases answering yes to sexual abuse were asked about sexual experiences; Categories were not mutually exclusive; A small minority of cases indicated additional sexual assaults occurring between the ages of 13 and 24 ( $n = 11$ ).

Emotional Maltreatment. Retrospective reports on emotional maltreatment from parents or guardians were obtained utilizing single-items that describe different experiences of emotional abuse (see Table 4). The items asked whether respondents had experienced six types of emotional maltreatment and scores were rated on a two-point (yes/no) format.

Table 4 *Experience of emotional maltreatment from parents / guardians by sex and child protection status*

	Total	Male	Female	Non Child protection case	Child protection case
Total (N)	2980	1579	1401	2128	852
Weighted (N)	2980	1555	1425	2794	186
Addressed in humiliating (e.g. being called lazy, stupid, or useless manner by parents/guardians	13.2	12.7	13.8	12.1	31.0
Humiliated or degraded in public by parents/guardians	5.4	4.5	6.5	4.7	16.7
Threatened about getting thrown out of the home by parents/guardians	13.7	15.4	11.7	12.3	33.8
Threatened about violent punishment by parents/guardians	3.1	3.2	2.9	2.5	12.2
Parents/guardians have through their behaviour shown that you were unwanted, unloved, and worthless	4.9	4.1	5.8	3.9	20.4
Parents/guardians have criticized or bullied you constantly	2.9	2.0	3.9	2.3	12.6
Total (%)	5.2	4.2	6.1	4.2	19.6

*Note.* All percentage values are weighted; Total = Positive endorsement of at least three items

Physical Neglect. Retrospective reports on neglect from parents or guardians were obtained utilizing seven single-items that describe the different experiences of physical neglect (aged<12; see Table 5). The items asked whether respondents had experienced seven types of physical neglect and scores were rated on a two-point (yes/no) format.

Table 5 *Experience of physical neglect from parents / guardians by sex and child protection status*

	Total	Male	Female	Non Child protection case	Child protection case
Total (N)	2980	1579	1401	2128	852
Weighted (N)	2980	1555	1425	2794	186
Aged <12 you were expected to wash own clothes	2.9	1.8	4.2	2.4	10.9
Aged <12 you had to attend school in dirty clothes because there were no clean ones available	1.5	1.6	1.4	1.1	8.1
Aged <12 you were occasionally starved due to lack of food or no one available to prepare meals	1.2	1.0	1.4	0.7	8.9
Aged <12 you were responsible for own care when sick	6.0	6.2	5.7	5.5	13.3
Aged <12 had to call a doctor for yourself when ill	0.6	0.5	0.6	0.5	2.1
Often had to care for yourself due to parental alcohol or drug problems	3.9	2.9	4.9	2.9	19.3
Were often abandoned / deserted in the home for several days	1.3	1.4	1.1	0.9	6.9
Total (%)	7.8	6.4	9.4	6.5	27.0

*Note.* All percentage values are weighted

### *Recoding of the variables*

For the ANOVA, the variables were recoded so that the answers 'don't know', 'not known' and 'irrelevant' were taken out of the analyses. Some items needed to be further recoded into the right factor. The factor 'self-concept' was developed by further recoding: The positive self-concept items were recoded into negative self-concept items, and all items were then summarized. The item of support was divided into social support, teacher support and family support, by summarizing the sub-items associated with these support groups. The factor 'contact with parents' was made by summarizing the items 'contact with mother' and 'contact with father'. Finally, the factor 'connectedness with parents' was created by summarizing the variables 'relationship with parents' and 'contact with parents'.

For the logistic regression analyses, the factors were recoded into a dummy, with the rule 'any score equal to or higher than 1 = 1'. These recoded factors were then put into the logistic regression analysis.

## Analysis

SPSS 18.0 was used to analyze the data. First, chi-square analyses were used for determining the significance of the relation between the types of maltreatment and suicidal thoughts and suicide attempts. Next, more chi-square analyses were done to find out if there was a significant relation between the types of maltreatment and the factors 'negative self-concept', 'social support', 'teacher support', 'family support', 'relationship with parents' and 'present contact with parents'. Furthermore, these factors were put into a chi-square analysis with suicide, suicide attempts and suicidal thoughts. A correlation analysis was used to look at the consensus between the support factors ('social support'; 'teacher support'; 'family support'; 'relationship with parents'; 'present contact with parents'). ANOVA was performed in order to determine whether the scores significantly vary across the factors. Moreover, logistic regression analysis was used to determine the influence of the factors on the relationships between the types of maltreatment and suicide, suicide attempts, and suicidal thoughts.

## Results

The percentage of maltreated children reporting suicidal thoughts and/or attempts is 19.7%, and of non-maltreated children is 18.8%. Table 6 shows the percentages of suicide (total), suicidal thoughts and suicide attempts for the types of maltreatment.

Table 6 *Percentages of Suicidal thoughts, Suicide Attempts and Suicide (Total) in Maltreated Children*

	<b>N</b>	<b>Suicidal thoughts</b>	<b>Suicide Attempts</b>	<b>Suicide (total)</b>
<b>Child Maltreatment (All Types)</b>	600	12.7%	7.0%	19.7%
<b>Physical Abuse</b>	253	11.5%	7.5%	19.0%
<b>Sexual Abuse</b>	116	8.6%	1.7%	10.3%
<b>Emotional Maltreatment</b>	257	16.0%	7.8%	23.7%
<b>Physical Neglect</b>	369	13.0%	6.5%	19.5%
<b>Non Maltreatment</b>	2279	13.7%	5.2%	18.8%

First, a correlation analysis was done to look at the consensus between the support factors ('social support'; 'teacher support'; 'family support'; 'relationship with parents'; 'present present contact with parents'). Social support was the least correlated factor, only teacher support and relationship with parents were positively related with social support. All other support variables were positively correlated with each other. Though, negative relationship existed for teacher support with family support, relationship with parents and present contact with parents. The results are shown in Table 7.

Table 7 *Correlations factors of support*

	Social Support	Teacher Support	Family support	Relation with parents	Contact with parents
Social Support	1	.242**	-.028	.060**	.003
Teacher Support	.242**	1	-.129**	-.040*	-.060**
Family support	-.028	-.129**	1	.209**	.186**
Relation with Parents	.060**	-.040*	.209**	1	.196**
Contact with parents	.003	-.060**	.186**	.196**	1

\*\* . Correlation is significant at the 0.01 level (2-tailed).

\* . Correlation is significant at the 0.05 level (2-tailed).

### *Chi-Square Analyses*

Chi-square analyses were used to determine which factors are significantly associated with which type of maltreatment. The analysis showed that maltreatment is positively related to teacher support\* and negatively to family support\* and relationship with parents\*. However, present contact with parents\* had a negative relation with physical abuse. Sexual abuse is negatively related to suicide (total)\*\* and suicide attempts\*. Furthermore, chi-square analysis on emotional maltreatment showed that it is positively related to 'suicide total'\*, and negatively to family support\*. Teacher support\*\* was positively associated with physical neglect. A negative relationship with parents\* however, had a negative relation with physical neglect. All the chi-square values were higher than 3.3.

Moreover, suicide (total) was positively associated with negative self-concept\*\*\* and teacher support\*\*\*. Suicide (total) had a negative relationship with family support\*\*\*, relation with parents\*\*\* and present contact with parents\*\*\*. The chi-square values were all higher than 11,7.

Suicidal thoughts were positively related to the same variables: negative self-concept\*\*\* and teacher support\*. The suicide variable had a negative association with family support\*\*\*, relationship with parents\*\*\* and present contact with parents\*\*\*. The chi-square values were higher than 5.9.

The variable suicide attempts however, were associated with all the variables. Negative self-concept\*\*\* and teacher support\* had a positive relationship with suicide attempts. Social support\*\*, family support\*\*\*, relationship with parents\*\*\* and contact with parents\*\*\* were negatively associated with suicide attempts. All chi-square values were higher than 5.10.

\*\*\*. Correlation is significant at the 0.001 level (2-tailed).

\*\* . Correlation is significant at the 0.01 level (2-tailed).

\* . Correlation is significant at the 0.05 level (2-tailed).

## ANOVA

The ANOVA analyses showed no significant results. Therefore, logistic regression analyses were done.

### *Logistic Regression Analyses*

#### Child Maltreatment

Logistic regression analysis showed which factors influence the relation between maltreatment and suicidal thoughts, suicide attempts and suicide (total) (see Table 8). It revealed that 'negative self-concept', 'social support', 'teacher support', 'relationship with parents' and 'present contact with parents' reliably predict suicide (total) in persons who have been maltreated in their childhood. The odds ratio of child maltreatment for suicide (total) was highest in 'negative self-concept' (OR=4.65) and more modest in 'teacher support' (OR=2.31). Furthermore, the risk of suicide is decreased by 'social support' (OR=0.45), 'relationship with parents' (OR=0.34) and 'present contact with parents' (OR=0.26). A 'negative self-concept' is associated with a nearly 7-fold increased risk of suicidal thoughts. Additionally, 'present contact with parents' decreases the risk of suicidal thoughts (OR=0.34). For suicide attempts, only 'teacher support' (OR=2.67) increases the risk. 'Relationship with parents' (OR=0.22) and 'present contact with parents' (OR=0.39), decrease the number of reports of suicide attempts.

#### Physical Abuse

Regression analysis (see Table 9) showed that negative 'self-concept' (OR=4.57) and 'teacher support' (OR=2.31) notably increase the scores of suicide (total). 'Present contact with parents' (OR=0.35) however, protected against the number of reports of suicide (total). Looking at suicidal thoughts, the odds ratio was high for 'negative self-concept' (OR=7.07).

#### Sexual Abuse

For the results of the regression analyses on sexual abuse, see Table 10. For sexual abuse, the score on suicide (total) is increased with a 13-fold chance by a 'negative self-concept'. Furthermore, 'present contact with parents' (OR=0.12) protected against the risk of suicide (total). A 'negative self-concept' (OR=9.99) increased the number of reports of suicidal thoughts.

Table 8 *Logistic Regression Analyses of Child Maltreatment*

	Suicidal Thoughts						Suicide Attempts						Suicide (Total)					
	B	S.E.	df	OR	95% C.I.	Sig.	B	S.E.	df	OR	95% C.I.	Sig.	B	S.E.	df	OR	95% C.I.	Sig.
	Neg. Self-concept	1,91	0,45	1,00	6,75	2,81-16,23	0,00	0,48	0,44	1,00	1,61	0,68-3,83	0,28	1,54	0,33	1,00	4,65	2,43-8,89
Social Support	-0,57	0,30	1,00	0,57	0,31-1,02	0,06	-0,76	0,43	1,00	0,47	0,20-1,08	0,07	-0,79	0,27	1,00	0,45	0,27-0,77	0,00
Teacher Support	0,44	0,31	1,00	1,56	0,85-2,86	0,15	0,98	0,40	1,00	2,67	1,22-5,87	0,01	0,84	0,28	1,00	2,31	1,34-3,97	0,00
Family Support	-0,25	0,56	1,00	0,78	0,26-2,33	0,66	-0,63	0,61	1,00	0,53	0,16-1,76	0,30	-0,78	0,52	1,00	0,46	0,17-1,27	0,13
Relation Parents	-0,50	0,30	1,00	0,61	0,34-1,11	0,10	-1,53	0,40	1,00	0,22	0,10-0,47	0,00	-1,08	0,26	1,00	0,34	0,20-0,57	0,00
Contact Parents	-1,08	0,34	1,00	0,34	0,17-0,67	0,00	-0,95	0,43	1,00	0,39	0,17-0,89	0,03	-1,37	0,31	1,00	0,26	0,14-0,47	0,00

Table 9 *Logistic Regression Analyses of Physical Abuse*

	Suicidal Thoughts						Suicide Attempts						Suicide (Total)					
	B	S.E.	df	OR	95% C.I.	Sig.	B	S.E.	df	OR	95% C.I.	Sig.	B	S.E.	df	OR	95% C.I.	Sig.
	Neg. Self-concept	1,96	0,76	1,00	7,07	1,61-31,18	0,01	0,64	0,68	1,00	1,89	0,50-7,18	0,35	1,52	0,51	1,00	4,57	1,67-12,51
Social Support	-0,36	0,48	1,00	0,70	0,27-1,79	0,45	-0,74	0,65	1,00	0,48	0,13-1,70	0,25	-0,59	0,42	1,00	0,55	0,24-1,25	0,16
Teacher Support	0,46	0,50	1,00	1,58	0,59-4,23	0,36	1,00	0,61	1,00	2,73	0,83-8,99	0,10	0,84	0,43	1,00	2,31	1,00-5,30	0,05
Family Support	-0,29	0,94	1,00	0,75	0,12-4,67	0,75	-1,10	0,93	1,00	0,33	0,05-2,05	0,24	-1,16	0,81	1,00	0,31	0,07-1,52	0,15
Relation Parents	-0,04	0,53	1,00	0,96	0,34-2,73	0,94	-1,00	0,59	1,00	0,37	0,12-1,18	0,09	-0,57	0,43	1,00	0,57	0,25-1,31	0,18
Contact Parents	-0,85	0,52	1,00	0,43	0,15-1,18	0,10	-0,78	0,63	1,00	0,46	0,13-1,58	0,22	-1,06	0,46	1,00	0,35	0,14-0,85	0,02

Table 10 *Logistic Regression Analyses of Sexual Abuse*

	Suicidal Thoughts						Suicide Attempts						Suicide (Total)					
	B	S.E.	df	OR	95% C.I.	Sig.	B	S.E.	df	OR	95% C.I.	Sig.	B	S.E.	df	OR	95% C.I.	Sig.
	Neg. Self-concept	2,30	1,16	1,00	9,99	1,02-97,91	0,05	1,04	#####	1,00	2,83	0,00	1,00	2,59	1,21	1,00	13,38	1,24-144,26
Social Support	-0,43	0,79	1,00	0,65	0,14-3,05	0,59	-0,69	#####	1,00	0,50	0,00	1,00	-0,54	0,78	1,00	0,58	0,13-2,68	0,49
Teacher Support	0,09	0,89	1,00	1,09	0,19-6,28	0,92	-0,38	#####	1,00	0,69	0,00	1,00	-0,01	0,91	1,00	0,99	0,17-5,89	0,99
Family Support	-2,24	1,24	1,00	0,11	0,01-1,20	0,07	18,50		1,00		0,00	1,00	-1,50	1,21	1,00	0,22	0,02-2,40	0,22
Relation Parents	0,24	0,90	1,00	1,27	0,22-7,40	0,79	-18,27	#####	1,00	0,00	0,00	1,00	-0,54	0,80	1,00	0,59	0,12-2,79	0,50
Contact Parents	-1,18	0,86	1,00	0,31	0,06-1,66	0,17	-19,28	#####	1,00	0,00	0,00	1,00	-2,09	0,80	1,00	0,12	0,03-0,59	0,01

### Emotional Maltreatment

The regression analysis (see table 11) showed that a 'negative self-concept' (OR=4.59) notably increases the risk of 'suicide total' in the sample. 'Social support' (OR=0.41), 'relationship with parents' (OR=0.31) and 'present contact with parents' (OR=0.36) decrease the score of suicide (total). Furthermore, 'negative self-concept' (OR=4.94) increases the risk of suicidal thoughts. For suicide attempts however, family support (OR=0.13) acts as a protective factor.

### Physical Neglect

The regression analysis (see table 12) revealed that 'negative self-concept' (OR=4.79) and 'teacher support' (OR=2.56) increase the risk of suicide (total) in this sample. 'Relationship with parents' (OR=0.36) and 'present contact with parents' (OR=0.35) however, protect against the risk. A 'negative self-concept' is associated with a 4-fold increased score of suicidal thoughts in persons who have been physically neglected in their childhood. Additionally, 'present contact with parents' decrease the number of reports of suicidal thoughts (OR=0.40). For suicide attempts, 'teacher support' (OR=3.00) increases the number of reports. A good 'relationship with parents' (OR=0.17), again, decreases the risk of suicide attempts.

Table 11 *Logistic Regression Analyses of Emotional Maltreatment*

	Suicidal Thoughts						Suicide Attempts						Suicide (Total)					
	B	S.E.	df	OR	95% C.I.	Sig.	B	S.E.	df	OR	95% C.I.	Sig.	B	S.E.	df	OR	95% C.I.	Sig.
Neg. Self-concept	1,60	0,56	1,00	4,94	1,64-14,93	0,01	0,79	0,70	1,00	2,20	0,56-8,71	0,26	1,52	0,46	1,00	4,59	1,85-11,40	0,00
Social Support	-0,37	0,43	1,00	0,69	0,34-10,16	0,39	-1,47	0,74	1,00	0,23	0,05-0,97	0,05	-0,89	0,41	1,00	0,41	0,19-0,91	0,03
Teacher Support	0,21	0,49	1,00	1,24	0,30-1,60	0,66	0,84	0,63	1,00	2,32	0,68-7,98	0,18	0,67	0,44	1,00	1,95	0,82-4,63	0,13
Family Support	0,61	0,87	1,00	1,84	0,48-3,21	0,48	-2,05	0,85	1,00	0,13	0,02-0,68	0,02	-0,97	0,71	1,00	0,38	0,10-1,51	0,17
Relation Parents	-0,79	0,44	1,00	0,45	0,19-1,07	0,07	-1,06	0,60	1,00	0,35	0,11-1,12	0,08	-1,16	0,40	1,00	0,31	0,14-0,68	0,00
Contact Parents	-0,58	0,53	1,00	0,56	0,20-1,57	0,27	-0,93	0,66	1,00	0,40	0,11-1,43	0,16	-1,02	0,49	1,00	0,36	0,14-0,94	0,04

Table 12 *Logistic Regression Analyses of Physical Neglect*

	Suicidal Thoughts						Suicide Attempts						Suicide (Total)					
	B	S.E.	df	OR	95% C.I.	Sig.	B	S.E.	df	OR	95% C.I.	Sig.	B	S.E.	df	OR	95% C.I.	Sig.
Neg. Self-concept	1,49	0,51	1,00	4,42	1,64-11,92	0,00	1,16	0,66	1,00	3,19	0,87-11,70	0,08	1,57	0,43	1,00	4,79	2,08-11,05	0,00
Social Support	0,07	0,37	1,00	1,07	0,52-2,21	0,85	-0,54	0,54	1,00	0,58	0,20-1,68	0,32	-0,16	0,33	1,00	0,85	0,45-1,62	0,62
Teacher Support	0,59	0,39	1,00	1,81	0,85-3,85	0,12	1,10	0,52	1,00	3,00	1,09-8,23	0,03	0,94	0,34	1,00	2,56	1,31-5,01	0,01
Family Support	-0,26	0,85	1,00	0,77	0,15-4,08	0,76	0,52	1,15	1,00	1,68	0,18-16,10	0,65	0,09	0,78	1,00	1,10	0,24-5,06	0,91
Relation Parents	-0,32	0,41	1,00	0,73	0,33-1,62	0,44	-1,75	0,53	1,00	0,17	0,06-0,49	0,00	-1,04	0,34	1,00	0,36	0,18-0,70	0,00
Contact Parents	-0,92	0,47	1,00	0,40	0,16-0,99	0,05	-0,71	0,59	1,00	0,49	0,16-1,55	0,23	-1,05	0,42	1,00	0,35	0,16-0,79	0,01

## Discussion

The relation between childhood maltreatment and suicide has been investigated. The focus was on this relationship and the influence of some factors on it. In this research, child maltreatment (all types), physical abuse, sexual abuse, emotional maltreatment and physical neglect ( $N=2879$ ) have been compared with each other on the subject of suicidal thoughts, suicide attempts and both (suicide total). The factors that were included in this study are 'negative self-concept', 'social support', 'family support', 'teacher support', 'relationship with parents' and 'present contact with parents'. It was expected that a 'negative self-concept' increases the risk of suicide. On the contrary, it was assumed that 'social support', 'family support', 'teacher support', 'relationship with parents' and 'present present contact with parents' act as protective factors of suicide.

The first research question was: '*Does childhood maltreatment influence the risk of suicidal thoughts and suicide attempts, and is there a difference for the different types of maltreatment?*' Chi-square analyses showed that maltreatment (all types) appeared not to be related to any of the suicide factors. Thus, the first hypothesis can be rejected. This is in contrast with much of the literature (Brown et al., 1999; Dube et al., 2001; Hahm et al., 2010; Meadows & Kaslow, 2002; Perroud et al., 2007; Sfoggia et al., 2008; Thompson et al., 2005). Furthermore, the assumption that has been made in this study that the interpersonal-psychological theory (Ribeiro & Joiner, 2009) can explain the death wish in maltreated persons, can't be confirmed because of these results. This indicates that, in general, the theory might not be applicable to childhood maltreatment. Possibly the items were too incoherent to measure one common factor. Therefore, for the second hypothesis it was investigated whether or not the several types of maltreatment differed in suicide outcome. The chi-square analyses revealed that physical abuse and physical neglect were not associated with the suicide factors. Emotional maltreatment did have a positive relation with suicide (total). However, sexual abuse gave the most unexpected results. It was negatively related to suicide attempts and suicide (total), which means that sexual abuse protects against suicidal risk behavior. This protective relation is in contrast with the studies of Bruffaerts et al. (2010) and Perroud et al. (2007), and difficult to explain. Perhaps sexual abuse causes more distress than the other maltreatment types, and therefore the victims of this type of maltreatment receive earlier and more often psychological help after the abuse has taken place. However, the results lead to the acceptance of the second hypothesis: there is a difference in the reports of suicidal thoughts and attempts for the four types of maltreatment. This seems to confirm the idea that the items were too unconnected to represent one joint factor. It indicates that in research, maltreatment should also be divided into the different types of maltreatment to obtain a more complete view on the subject.

Finally, the research question was proposed: '*Which factors influence the relation*

*between childhood maltreatment and suicide?’* In order to answer this question, a correlation analysis and a logistic regression analysis was been done. First, the correlation analysis was executed for the support variables to look at the cohesion between these variables. It revealed that social support was the least correlated factor, but the other support variables were all correlated with each other. Social support was positively related with teacher support and relationship with parents, the rest of the support variables were positively correlated with each other. However, teacher support was negatively correlated with family support, relationship with parents and present contact with parents. It indicates that maltreated persons who have received support from teachers, received less support from their family, had a worse relationship with their parents and have less contact with them at the moment. In conclusion, maltreated persons received either support from their family and/or parents, or they received support from their teacher(s). Furthermore, the correlation analyses suggest that the support variables do not represent one common factor, which implicates that the results of the regression analyses won't be similar for these variables. For the following logistic regression analysis, the interpretations of the results are arranged per factor.

#### *Negative Self-concept*

In the current research, the factor 'negative self-concept' was the most important factor for the relation between maltreatment and suicide. Negative self-concept has a significant influence on the relation between all types of childhood maltreatment and suicidal thoughts and suicide (total). This means that a high score on 'negative self-concept' leads to a higher score on suicide. Given the high OR values in every type of maltreatment, 'negative self-concept' has a considerably high impact. That is, the hypotheses for suicidal thoughts and suicide (total) can fully be accepted for all the types of maltreatment. This is in line with the study of Eskin et al. (2007) and the assumption concerning the interpersonal-psychological theory (Ribeiro & Joiner, 2009). The concordance with the theory indicates that the misinterpretation that the self is incompetent, leads to the increased risk of suicide ideation in maltreated persons. This means that, in supporting the theory, maltreatment has an additive value. In addition, the results for sexual abuse are remarkable, because it's in contrast with the results of the chi-square analyses. Persons who have been sexually abused have a lower risk for suicidal risk behavior, but sexually abused person with a negative self-concept have a higher risk for suicidal risk behavior. This contradiction is difficult to elucidate. Furthermore, the hypothesis must be rejected for suicide attempts, since there wasn't any significance between 'negative self-concept' and this outcome variable. Perhaps it could be explained by the nature of negative self-concept. Persons with a negative self-concept have negative thoughts about themselves. The transition from negative thoughts about yourself to suicidal thoughts seems smaller for maltreated persons than the transition to suicide attempts, which might explain the differences in the results. Therefore, more research is

needed to examine the relationships between a negative self-concept, suicidal thoughts and suicide attempts in persons who have experienced any form of childhood maltreatment.

#### *Social support*

Kaslow et al. (2002) found that social support (from family and friends) significantly protects against attempting suicide in low-income African American women. In the chi-square of the current research as well, social support was significantly related to suicide attempts. However, it was not related to suicidal thoughts, suicide (total) and any of the maltreatment types. The logistic regression analyses showed that social support is a protector of suicide (total) in the conditions of combined maltreatment and emotional maltreatment. That is, the hypotheses on social support can be accepted for suicide (total) in these two conditions. Perhaps, the results might be more conclusive if the item also takes the amount of received social support into account.

#### *Teacher support*

The factor 'teacher support' gave some unexpected results; all hypotheses of teacher support must be rejected. It was expected that teacher support is a protective factor of suicide. This assumption was based on the studies of Eisenberg et al. (2007) and Fleming et al. (2007). These researches were limitedly applicable to the current study, because the first one focused solely on students who were sexually abused, and the second study did not include childhood maltreatment at all. The logistic regression showed that participants in the combined maltreatment condition and the physical neglect condition, scored higher on 'suicide attempts' and 'suicide total' when they received more teacher support. Participants in the physical abuse condition also had significantly higher scores on suicide (total). That means that teacher support is a risk factor of suicide in these conditions. The fact that teacher support was a risk factor for suicidal behavior could be explained by the assumption that the participants only received teacher support when they were in negative circumstances. Moreover, another indication is that teacher support did not help the child sufficiently. This would mean that improvement is needed in teacher support in Denmark.

#### *Family support*

Literature showed that family support is a protective factor of suicidal behaviors (Au et al., 2009; Randell et al., 2006; Sharaf et al., 2009). In this comprehensive research, the maltreatment types were also included in the logistic regression analyses. Participants who have been emotionally maltreated and received family support, had lower scores on the variable suicide attempts. Thus, the hypothesis for suicide attempts can solely be accepted for the emotional maltreatment condition. All the other hypotheses of family support must be rejected, because of a lack of significant relations. These results were unexpected, and difficult to elucidate. The results might be more plausible if the item would also ask about the amount of received family support. More research is needed on the subject of family support

with childhood maltreatment and suicidal behavior to give a clear explanation of this outcome.

#### *Relationship with Parents*

A good relationship with parents acted as a protective factor of suicide attempts and suicide (total) in the combined maltreatment condition. It means that the hypotheses for suicide attempts and suicide (total) can be accepted for maltreatment (all types). This outcome is coherent with the literature (Eisenberg et al., 2007; Salzinger et al., 2007). Furthermore, the assumption that the interpersonal-psychological theory can be adopted in the association between maltreatment and suicide can't be fully confirmed, because of a lack of significance for suicidal thoughts (Ribeiro & Joiner, 2009). For physically neglected participants as well, scores on suicide attempts and suicide (total) were reduced by a good relationship with parents. Therefore, the hypotheses for suicide attempts and suicide (total) can also be accepted for physical neglect. Moreover, emotionally maltreated persons scored lower on suicide (total), when they had a good relationship with their parents. Thus, the hypothesis for suicide (total) can also be accepted for the emotional maltreatment condition. The hypotheses must be rejected for physical abuse and sexual abuse. This is in contrast with the previously mentioned researches. The contradiction might be explained by the characteristics of these studies. The study of Aspelmeier et al. (2007) was aimed solely on sexually abused female students. Additionally, Salzinger et al. (2007) concentrated on preadolescents with the age of 9-12 years old, and used a considerably smaller sample size. The current study is more representative, since a large sample size ( $N=2980$ ) was used that consisted of almost equal numbers of 24 year old males and females.

#### *Present contact with parents*

The factor 'present contact with parents' has not yet been included in research on the association between maltreatment and suicidal behavior. In this research, logistic regression analyses showed that present contact with parents was – in relationship with all the suicide variables – of significant influence in maltreatment (all types). This leads to the acceptance of the hypotheses for all the suicide variables in this condition. Moreover, the assumption that the interpersonal-psychological theory (Ribeiro & Joiner, 2009) can be applicable on the relationship between childhood maltreatment and suicide is also confirmed by these results. For the physical neglect condition, present contact with parents significantly reduced the scores on suicidal thoughts and suicide (total). The herewith connected hypotheses can be accepted. Additionally, participants in the conditions of physical abuse, sexual abuse, and emotional maltreatment also had lower scores on suicide (total) if they had more contact with their parents. That is, the relationship between all the maltreatment types and suicide (total) were negatively influenced by the factor 'present contact with parents'. It seems important

that contact with parents is being stimulated, in order to protect against suicidal behavior in maltreated persons.

### *Conclusion*

Maltreatment (all types) had no significant relationship with suicidal risk behavior. Additionally, some differences have been found for the different types of maltreatment. Sexual abuse did have a relation with suicide attempts and suicide (total), and emotional maltreatment with suicide (total) as well. All the other relationships between the maltreatment types and the suicide variables were not significant. The chi-square analyses of the factors showed significance between the suicide variables and almost all factors. However, there was hardly any significance between the factors and the maltreatment types. In the logistic regression analyses, the factors did influence the association between maltreatment and suicide in different ways. The most important factor was 'negative self-concept', followed by 'present contact with parents'. Teacher support gave the most remarkable result: it had the status of a risk factor instead of a protective factor of suicidal risk behavior.

### *Strengths and Limitations*

In this study, the sample size was larger than most samples in research that has been done concerning the relation between maltreatment and suicide. Furthermore, the interviews were done by trained professionals. Moreover, the submission of the Child Protection Cases in this study results in a more valid comparison of maltreated persons and non-maltreated persons. Additionally, because of the similarities in age of the participants, it's more reliable to make comparisons within the sample. Another added value offered by this research concerns the fact that all types of maltreatment were taken into account as well as one combined factor. This led to an extensive view on the subject. Furthermore, many factors have been included in this study, which also contributes to an extensive view on the subject. Finally, the factor 'present contact with parents' provides some new insights on the relation. 'Present contact with parents' was not yet taken into account in earlier researches, which makes this factor and the research innovative.

Several limitations of this research should be noted as well. It's difficult to generalize the results and implications to other age groups in the population, because of the similar age of the participants. Furthermore, the information from the variables could have been restricted, since the variables were made dichotomous for the analyses. Finally, another limitation of this research is that it uses self-reports. However, the research keeps its strengths and implications. The largest and most representative research that had been done in 21 countries, did not include all types of maltreatment (Bruffaerts et al., 2010). Therefore, it is recommendable that a nationwide research like this one should be done in all countries. In the Netherlands, the home country of the authors, few research has been done concerning

the association between all the types of childhood maltreatment and suicidal risk behavior, thus it's specifically recommended for the Netherlands.

## References

- Affifi, T.O., Boman, J., Fleisher, W., & Saree, J. (2009). The relationship between child abuse, parental divorce, and lifetime mental disorders and suicidality in a nationally representative adult sample. *Child Abuse & Neglect*, *33*, 139–147.
- Arata, C.M., Langhinrichsen-Rohling, J., Bowers, D., & O' Brien, N. (2007). Differential correlates of multi-type maltreatment among urban youth. *Child Abuse & Neglect*, *31*, 393–415.
- Aspelmeier, J.E., Elliott, A.N., & Smith, C.H. (2007). Childhood sexual abuse, attachment, and trauma symptoms in college females: The moderating role of attachment. *Child Abuse & Neglect*, *31*, 549-566.
- Au, A.C.Y., Lau, S., & Lee, M.T.Y. (2009). Suicide ideation and depression: The moderation effects of family cohesion and social self-concept. *Adolescence*, *44*(176), 851-868.
- Boxer, P. (2010). Variations in Risk and Treatment Factors Among Adolescents Engaging in Different Types of Deliberate Self-Harm in an Inpatient Sample. *Journal of Clinical Child & Adolescent Psychology*, *39*(4), 470–480.
- Brown, J., Cohen, P., Johnson, J.G., & Smailes, E.M. (1999). Childhood Abuse and Neglect: Specificity of Effects on Adolescent and Young Adult Depression and Suicidality. *Journal of the American Academy of Child and Adolescent Psychiatry*, *38*(12), 1470-1496.
- Bruffaerts, R., Demyttenaere, K., Borges, G., Haro, J.M., Chiu, W.T., Hwang, I., ... Nock, M.K. (2010). Childhood adversities as risk factors for onset and persistence of suicidal behavior. *The British Journal of Psychiatry*, *197*, 20-27.
- Burns, E.E., Jackson, J.L., & Harding, H.G. (2010). Child Maltreatment, Emotion Regulation, and Posttraumatic Stress: The Impact of Emotional abuse. *Journal of Aggression, Maltreatment & Trauma*, *19*(8), 801-819.
- Cantón-Cortés, D., & Cantón, J. (2010). Coping with child sexual abuse among college students and post-traumatic stress disorder: The role of continuity of abuse and relationship with the perpetrator. *Child Abuse & Neglect*, *34*(7), 496-506.
- Cheng, Y., Tao, M., Riley, L., Kann, L., Ye, L., Tian, X., Tian, B., Hu, J., & Chen, B. (2009). Protective factors relating to decreased risks of adolescent suicidal behaviour. *Child: care, health and development*, *35*(3), 313–322.
- Cicchetti, D., & Toth, S.L. (2005). Child Maltreatment. *Annual Reviews*, *1*, 409-438
- Compton, M.T., Thompson, N.J., & Kaslow, N.J. (2005). Social environment factors associated with suicide attempt among low-income African Americans: The protective

- role of family relationships and social support. *Social Psychiatry and Psychiatric Epidemiology*, 40(3), 175-185.
- Dervic, K., Oquendo, M.A., Grunebaum, M.F. Ellis, S., Burke, A.K., & Mann, J.J. (2004). Religious Affiliation and Suicide Attempt. *The American Journal of Psychiatry*, 161(12), 2303-2308.
- Ditlevsen, D.N., & Elklit, A. (2010). The combined effect of gender and age on post traumatic stress disorder: do men and women show differences in the lifespan distribution of the disorder? *Annals of General Psychiatry*, 9(32).
- Dube, S.R., Anda, R.F., Felitti, V.J., Chapman, D.P., Williamson, D.F., & Giles W.H. (2001). Childhood Abuse, Household Dysfunction, and the Risk of Attempted Suicide Throughout the Life Span. Findings From the Adverse Childhood Experiences Study. *American Medical Association*, 286(24), 3089-3096.
- Eisenberg, M.E., Ackard, D.M., & Resnick, M.D. (2007). Protective Factors and Suicide Risk in Adolescents with a History of Sexual Abuse. *Journal of Pediatric Psychology*, 151, 482-7.
- Eskin, M., Ertekin, K., Dereboy, C., & Demirkiran, F. (2007). Risk Factors for and Protective Factors Against Adolescent Suicidal Behavior in Turkey. *Crisis*, 28(3), 131–139.
- Fleming, T.M., Merry, S.N., Robinson, E.M., Denny, S.J., & Watson P.D. (2007). Self-reported suicide attempts and associated risk and protective factors among secondary school students in New Zealand. *Australian and New Zealand Journal of Psychiatry*, 41(3), 213-221.
- Hahm, C.H., Lee, Y., Ozonoff, A., & Van Wert, M.J. (2010). The Impact of Multiple Types of Child Maltreatment on Subsequent Risk Behaviors Among Women During the Transition from Adolescence to Young Adulthood. *Journal of Youth and Adolescence*, 39, 528-540.
- Kaslow, N.J., Thompson, M.P., Okun, A., Price, A., Young, S., Bender, M., Wyckoff, S., Rollins, E., Twomey, H., Goldin, J., & Parker, R. (2002). Risk and Protective Factors for Suicidal Behavior in Abused African American Women. *Journal of Consulting and Clinical Psychology*, 70(2), 311–319.
- Kim, J., & Cicchetti, D. (2010). Longitudinal pathways linking child maltreatment, emotion regulation, peer relations, and psychopathology. *Journal of Child Psychology and Psychiatry*, 51(6), 706-716.
- Locke, T.F., & Newcombe, M.D. (2005). Psychosocial Predictors and Correlates of Suicidality in Teenage Latino Males. *Hispanic Journal of Behavioral Sciences*, 27(3), 319-336.
- May-Chahal, C., & Cawson, P. (2005). Measuring child maltreatment in the United Kingdom:

- A study of the prevalence of child abuse and neglect. *Child Abuse & Neglect*, 29(9), 969-984.
- Meadows, L.A., & Kaslow, N.J. (2002). Hopelessness as Mediator of the Link Between Reports of a History of Child Maltreatment and Suicidality in African American Women. *Cognitive Therapy and Research*, 26(5), 657–674.
- Perroud, N., Courtet, P., Vincze, I., Jaussent, I., Jollant, F., Bellivier, F., Leboyer, M., Baud, P., Buresi, C., & Malafosse, A. (2007). Interaction between BDNF Val66Met and childhood trauma on adult's violent suicide attempt. *Genes, Brain and Behavior*, 7, 314–322.
- Rachman, S. (2004). *Anxiety, Second Edition*. New York: Psychology Press.
- Randell, B.P., Wang, W., Herting, J.R., & Eggert, L.L. (2006). Family Factors Predicting Categories of Suicide Risk. *Journal of Child and Family Studies*, 15(3), 255-270.
- Ribeiro, J.D., & Joiner, T.E. (2009). The Interpersonal-Psychological Theory of Suicidal Behavior: Current Status and Future Directions. *Journal of Clinical Psychology*, 65(12), 1291-1299.
- Ryan, K.D., Kilmer, R.P., Cauce, A.M., Watanabe, H., & Hoyt, D.R. (2000). Psychological Consequences of Child Maltreatment in Homeless Adolescents: Untangling the Unique Effects of Maltreatment and Family Environment. *Child Abuse & Neglect*, 24(3), 333–352.
- Salzinger, S., Rosario, M., Feldman, R.S., & Ng-Mak, D.S. (2007). Adolescent Suicidal Behavior: Associations With Preadolescent Physical Abuse and Selected Risk and Protective Factors. *Journal of the American Academy of Child and Adolescent Psychiatry*, 46(7), 859-866.
- Schotte, D.E., & Clum, G.A. (1982). Suicide Ideation In A College Population: A Test Of A Model. *Journal of Consulting and Clinical Psychology*, 50(5), 690-696.
- Scott, K.M., Smith, D.R., & Ellis, P.M. (2010). Prospectively Ascertained Child Maltreatment and Its Association With DSM-IV Mental Disorders in Young Adults. *Archives of General Psychiatry*, 67(7), 712-719.
- Senn, T.E., & Carey, M.P. (2010). Child maltreatment and women's adult sexual risk behavior: Childhood sexual abuse as a unique risk factor. *Child Maltreatment*, 15(4), 324-335.
- Sfoggia, A. Pacheco, M.A., & Grassi-Oliveira, R. (2008). History of Childhood Abuse and Neglect and Suicidal Behavior at Hospital Admission. *Crisis*, 29(3), 154–158.
- Sharaf, A.Y., Thompson, E.A., & Walsh, E. (2009). Protective Effects of Self-Esteem and Family Support on Suicide Risk Behaviors among At-Risk Adolescents. *Journal of Child and Adolescent Psychiatric Nursing*, 22(3), 160-168.
- Simpson, J.A., & Rholes, W.S. (1998). *Attachment Theory and Close Relationships*. New

York: The Guilford Press.

Thompson, R., Briggs, E., English, D.J., Dubowitz, H., Lee, L.C., Brody, K., Everson, M.D., & Hunter, W.M. (2005). Suicidal Ideation Among 8-Year-Olds Who Are Maltreated and At Risk: Findings From the LONGSCAN Studies. *Child Maltreatment*, 10(1), 26-36.

World Health Organization (2010, August). *Health topics. Fact sheets: Child Maltreatment*. Retrieved May 11, 2011, from

<http://www.who.int/mediacentre/factsheets/fs150/en/index.html>

## Appendix

### SFI – OVERSIGHT

Questions	(SPSS variable name)	Variables
1	<b>(5) Gender</b>	Man: 1, woman: 5, not known: 9
2	<b>(6) Living accommodations</b>	8 options
3	<b>(7) Number of children living at home</b>	0: 0 to 7:7 children, 8: don't know, 9: not known.
4	<b>(8) Education level</b>	9 options, of which one is "don't know"
5	<b>(9) Ever started on vocational or higher education?</b>	5 options: 1: completion, 2: under education, 3: dropped out 5: never started on one, 9: not known.
6	<b>(10) Type of vocational or higher education.</b>	8 options, of which one is no education.
7	<b>(11) Main occupation</b>	17 options.
8	<b>(12) Weekly working hours</b>	Write 888, if you don't know.
9	<b>(13) civil status – married or living together</b>	Options: 1: married or registered partnership, 2: living together, 5: no, 8: don't know, 9: not known.
10	<b>(14) Boy/girlfriend – going steady</b>	Options: 1: yes, 5: no, 8: don't know, 9: not known, 10: irrelevant.
11	<b>(15) Social support from partner during problems</b>	5 options: 0: haven't had any problems, 1: always, 2: usually, 5: no/rarely, 8: don't know, 9: not known, 10: irrelevant.

- 12 (16-30) Questions about selfconception**  
 15 items with 4 point Likert scale from 1: high agreement to 4: high disagreement, 8: don't know, 9: not known.  
 A: selfsatisfaction, b: can't do anything, c: good qualities, d: able to do things as good as others, e: have a lot to be proud of, f: feels useless, g: feel as valuable as others, h: want more self respect, i: tendency to feel like a failure, j: positive self attitude, k: need a true friend, l: nobody to confess inner thoughts too, m: often feels empty, n: often feels rejected by others, o: often lack self confidence.
- 13 (31-34) Selfrating of physical and mental health.**  
 Options 1: yes, 5: no, 8: don't know, 9: not known.  
 A: disturbance in sleep, b: anxiety attacks, c: usage of nerve medicine or sleeping pills, d: received help for mental problems from psychologist, psychiatrist, therapy group or at hospital.
- 14 (35) The presence of abdominal pain during the last 3 months**  
 6 options: 1: daily, 2: weekly, 3: each month, 4: not at all/rare, 8: don't know, 9: not known.
- 15 (36) Headache during the last 3 months**  
 Options like question 14.
- 16 (37) Feelings of overwhelming the last 3 months**  
 Options like question 14
- 17 (38) Nervous and unstable the last 3 months**  
 Options like question 14.
- 18 (39) Feeling down and sad the last 3 months**  
 Options like question 14, with extra option for not wanting to participate: 7.
- 19 (40) Irritability for no reason the last 3 months**  
 Options like question 18.
- 20 (41) Feelings of problems being overwhelming the last 3 months**  
 Options like question 18.

- 21 (42-45) recollections of past unpleasant experiences**  
Options: 1: yes, 5: no, 8: don't know and 9: not known to the following categories:  
A: nightmares/unwillingly thinking, b: deliberate avoidance of situation/recollection, c: constant hyper vigilance, d: emotional detachment from others and surroundings.
- 22 (46-55) statements about personality**  
Options 3 point likert from: 1: don't fit, 2: partly fit, 3: good fit, 8: don't know, 9: not known in the following categories:  
A: easily angry and often hot-tempered, b: keeps to yourself/loner, c: obedient, d: have at least one good friend, e: often starts conflicts and ability to make others do things, f: peers are mostly like you, g: often accused of lying and stealing h: often teased or bullied, i: steals, j: trouble getting friends.
- 23 (56-61) Statements about feelings and behaviour the last 6 months.**  
Options: 5 point likert, 1= never – 5= very often, 8: don't know, 9: not known in the following categories:  
A: problems with finishing projects, b: difficulty with handling demanding assignments, c: problems with remembering appointments or other things, which ought to be remembered, d: avoidance or procrastination of assignments, which demands considerations, e: restlessness in hands and feet, when required to sit still, f: feelings of hyperactivity and need to do things, because of an inner drive.
- 24 (62) Health selfrating**  
5 point Likert: 1: very good to 5: very bad, 8: don't know, 9: not known.
- 25 (63) Reduced working ability because of sickness, accident, handicap or hard work**  
3 point Likert: 1: yes, very much, 3: no, not very much, 5: no, not at all., 8: don't know, 9: not known.
- 26 (64) "Has a doctor told you, what is wrong with you?"**  
1: yes, 5: no, 8: don't know, 9: not known, 10: irrelevant.
- 27 (65) Chronic disease or disorders**  
1: yes, 5: no, 8: don't know, 9: not known.
- 28 (66) Current mental disorder**  
1: yes, 5: no, 8: don't know, 9: not known.

- 29 (67) “Has your doctor told you that you have a mental disorder?”**  
3 point Likert: 1: yes, very much, 3: no, not very much, 5: no, not at all, 8: don't know, 9: not known.
- 30 (68) Alcohol problems**  
Options: 1: yes, but not anymore, 2: yes, 5: no, never, 8: don't know, 9: not known.
- 31 (69-86) Alcohol consumption on the latest weekday (not Friday, Saturday or Sunday)**  
Options: number of bottles, number of glasses, and number of units of alcohol in the following categories (no coding in SPSS).  
1. Regular beer, 2. Ready to drink (Barcardi breezer etc.), 3. strong beer, 4. Red/white wine, 5. Dessert wine, 6. liquor.
- 32 (87) Height**  
Centimetres.
- 33 (88) Weight**  
Kilograms.
- 34 (89-96) Drugs “Have you ever tried...?”**  
Options: 1: Yes, 5: no, 8: don't know, 9: not known in the categories:  
A: speed, b: cannabis, c: cocaine, d: LSD, e: euphoriant mushrooms, f: heroine, g: “sniffing”, h: other drugs.
- 35 (97) Age of drug debut**  
Age in years.
- 36 (98) Last consumption of the above listed drugs**  
1: one week, 2: one month, 3: last 3 months, 4: last year, 5: more than one year, 8: don't remember/know, 9: not known.
- 37 (99) suicide attempts and –thoughts**  
4 Options: 1: yes, considerations, 2: yes, attempted, 5: no, 8: don't know, 9: not known.
- 38 (1001) Willingly overdose or other forms of self harming behaviour**  
5 options; 1: yes, overdose, 2: yes, self harm, 3: yes, both, 4: No to both, 5: don't know (no coding in SPSS).

**A: (101) The last time of overdose**

4 options: 1: less than a month, 2: between a month and a year, 3: more than a year, 8: don't know/remember, 9: not known, 10: irrelevant.

**B: (102-109) Any of the following reasons explaining overdose?**

5 options: 1: yes, 5: no, 7: don't want to answer, 8: don't know, 9: unknown, 10: irrelevant in the "categories":

A: wanting to show desperation, B: death wish, c: self punish, d: scare somebody, e: revenge, f: intolerable feelings, g: find out if anyone really cares, h: attention wish.

**C: (110) Hospitalization because of overdose?**

Options: 1: yes, 5: no, 8: don't know/remember, 9: not known, 10: irrelevant.

**39 A: (111) Time of last self harming**

4 options: 1: less than one month, 2: between one month and a year, 3: more than a year, 4: don't know (no SPSS coding).

**B: (112-119) any of the following reasons explaining self harm?**

Options: 1: yes, 5: no, 7: don't wish to answer, 8: don't know/remember, 9: not known, 10: irrelevant in the following categories

A: wanting to show desperation, B: death wish, c: self punish, d: scare somebody, e: revenge, f: intolerable feelings, g: find out if anyone really cares, h: attention wish.

**C: (120) Hospitalization because of selfharm?**

Options: 1: yes, 5: no, 8: don't know, 9: not known, 10: irrelevant.

**40 (121-128) Statements about body, weight and eating habits**

6 options with 4 point Likert after how good the statements fit: 1: very good fit – 4: never fit, 8: don't know, 9: unknown. Statements in following categories:

A: bad conscience, when eating candy, b: On diet, c: satisfied with eating habits, d: to avoid gaining weight or for loose of weight: starvation diet or faste, e: to avoid gaining weight or for loose of weight: diet pills or diet powder, f: to get rid of eat food: vomit, g: find it uncomfortably eating with others, h: afraid of not being able to stop eating, when first started.

**41 (129) Relationship to biological parents - childhood.**

4 options: 1: loving and friendly, 2: neutral, 3: conflict-ridden, 8: don't know, 9: not known.

**42 (341-343) Living, telephoning or texting with biological father and mother.**

1. Living with parents (SPSS codes: 1: yes, both, 2: yes, mother, 3: yes, father, 5: no, 7: both dead, 8: don't know, 9: not known) ,
2. Contact to mother (SPSS coding: 1: once or more each week, 2: rare, 3: never, 7: mother dead, 8: don't know, 9: not known, 10: irrelevant)
3. Contact to father (SPSS coding: 1: once or more each week, 2: rare, 3: never, 7: father dead, 8: don't know, 9: not known, 10: irrelevant)

**43 (131) Hit by parents/stepparents**

2 options: 1: yes, 5: no, 8: don't know, 9: not known.

**A (132): age the last time of beating**

5 options: 1: below the age of 5, 2: 5-6 years, 3: 7-9 years, 4: at least 10 years, 8: don't know, 9: not known, 10: irrelevant.

**B. (133) Age the first time of beating**

5 options: 1: 0-5 years, 2: 6-12 years, 3: 13-18 years, 4: 19 – years, 5: don't know (no SPSS coding).

**C. (134) repeated beatings over a longer period?**

5 options: 1: repeated times during several years, 2: repeated times over one period, 3: single times. 4: only once, 8: don't know, 9: not known, 10: irrelevant.

**44 (135-242) Events, which the subject, family or close friends have experienced.**

Options: 1: yourself, 2: family, 3: friends, 4: No one, SPSS coding is 1: answered, 5: unanswered, 8: don't no, 9: not known with 5 variables per event category:

A: serious traffic accident, b: fire, c: another accident, d: assault, e: threats about violence, f: drowning, g: robbery, h: maltreatment, i: bullying or humiliation, j: rape, k: threaten suicide, l: attempted suicide, m: committed suicide, n: died, o: a serious physical disease, p: mental disorder, q: eating disorders, r: repeated physical punishment from family member, s: sexual assault committed by family member, t: received help from psychologist, u: participated in a self help group, v: medical ordered medicine for emotional or mental problem.

- 45 (243-248) Have you (as an adult – over 18 years) experienced the following types of violence?**  
 4 options: 1: yes, within the last 12 months, 2: yes, 5: no, 8: don't know/remember, 9: unknown.
- Violence categories:  
 A: threats of physical harm, b: pushed, pulled or hit with a light touch, c: kicked, hit with fist or object, d: thrown into furniture, walls, down staircase, e: choking attempts, attacked with knife or firearm, f: other forms of violence.
- 46 (249) As an adult have you ever been subject of threats of violence that were to serious that you got scared?**  
 4 options: 1: yes, during the last 12 months, 2: yes: earlier, 5: no, 8: don't know, 9: not known.
- 47 (250-259) Who has threatened you or exposed you to violence?**  
 1 option for marking yes in the following categories, SPSS coding is 1: answered, 5: not answered, 8: don't know, 9: not known, 10: irrelevant:
- A: current spouse/partner, b: former spouse/partner, c: current or former boy-/girlfriend, d: parents or stepparents, e: other family members, f: friend or acquaintance, g: colleague or person on your work, h: personal on institution/school, i: another person, which I know, j: a stranger, k: don't know.
- 48 (260-265) forced to sexual activity**  
 6 options: 1: yes, as a child (below 13), 2: yes, as adolescent (13-17), 3: as an adult (18-), 4: yes, but don't remember when, 5: No, 6: unknown, SPSS coding is 1: answered, 5: not answered, 8: don't know, 9: not known.
- 49 (266-269) Was it...**  
 Options: 1: yes, 5: no, 8: don't know, 9: not known in categories:  
 a. Sexual touch, flasher, b: attempted forced intercourse, c: forced intercourse, d: sexual abuse.
- 50 (270) Has it happen in the last year?**  
 1: yes, 5: no, 8: don't know, 9: not known.

**51 (271-281) who forced you?**

1 option for marking yes to the following categories, SPSS coding is 1: answered, 5: not answered, 8: don't know, 9: not known:

a. Current spouse/partner, b: former spouse/partner, c: current or former girl-/boyfriend, d: parents/stepparents, e: other family members, f: a friend or acquaintance, g: colleague or person on your work, h: personal on institution/school, i: another person, which I know, j: a stranger, k: don't know.

**52 (282-290) treated in the following ways by parents/stepparents:**

3 options for marking 1: yes, 5: no, 8: don't know, 9: not known.

Categories:

A: spanking, b: beaten with objects, c: threaten by weapon, d: thrown things after you, e: choking, f: burn or bite marks, g: repeated bruises after beating, h: doctor pointed out injuries on you for example broken bones, burns etc. I: violence resulting in bruises, bleedings or other physical harm.

**53 (291) Told other adults outside the family about it**

Options: 1: yes, 5: no, 8: don't know, 9: not known.

**54 (292-294) Have you been examined after the incident by the following:**

Options: 1: yes, 5: no, 8: don't know, 9: not known.

Categories:

A: doctor, B: teacher, social worker, nurse or school teacher, c: other adults outside the family.

**55 (295-300) Experienced the following in childhood or youth.**

Options: 1: yes, 5: no, 8: don't know, 9: not known in the following categories:

A: Addressed in humiliating or degrading manner by parents/stepparents, b: humiliated or degraded in public by parents/stepparents, c: threatened about getting thrown out of the home by parents/stepparents, d: threatened about violent punishment by parents/stepparents, e: parents/stepparents have through their behaviour shown that you were not wanted or loved and worthless, f: parents/stepparents have criticized or bullied you.

The self report items ends here and the rest of the questions are ask by an interviewer

- 56 (301-307) Parents have different options about when a child should be able to take care of themselves – below 12 years did you experience the following:**  
Options: 1: yes, 5: no, 8: don't know, 9: not known in the following categories:
- A: you washed your own close, b: went to school in dirty clothes, c: make sure yourself that you went to the dentist, d: went hungry, because there was no one to cook you dinner or nothing in the fridge, e: had to take care of younger siblings, while parents were out, f: take care of yourself, while you were sick, g: had to call an doctor yourself, when sick.
- 57 (308-310) Responsibilities as a child, ever experienced:**  
Options: 1: yes, 5: no, 8: don't know, 9: not known in the following categories:
- A: take care of yourself, because your parents had problems with alcohol and drugs, b: you parents depended on you help, because they had emotional problems, c: often take care of yourself, because your parents went away or stayed away for several days at the time.
- 58 (311-316) How did you experience your childhood?**  
7 options: 7 point Likert scale from 1: never, 4:sometimes to 7: always, 8: don't know, 9: not known.
- How often...
- a: were you overall satisfied with support received as a child, b: did you have contact to others in the same situation, c: could you talk about your thoughts and feelings as a child, d: were there someone to help you with practical matters as a child, e: did you feel let down by people, whom you expected you could count on, f: were there someone, who helped you or supported you, when you needed it as a child.
- 59 (317-327) Who helped and supported you**  
1 option for marking yes, SPSS coding: 1: answered, 5: unanswered, 8: don't know, 9: not known, 10: irrelevant.
- A: mother, b: father, c: grandparents, d: siblings, e: other family, f: friends, g: friends' parents, h: school teacher, i: teachers, j: others
- 60 (328) seriously wanted to get away from home, before the age of 16.**  
Options, 1: yes, 5: no, 7: don't want to answer, 8: don't know/remember, 9: not known.

- 61      **(329) Ran away from home in 24 hours or more**  
Options: 1: yes, 5: no, 8: don't know/remember, 9: not known.
- 62      **(330-336) Committed one of following acts**  
3 options; 1: yes, 5: no, 8: don't know, 9: not known.
- Categories:  
A: shoplifting, b: stolen bike, c: stolen car, d: burglary, e: vandalism, f: violence, received a conviction for crime.
- 63      **(337) Number of schools in the first 9 years of school attendance.**
- 64      **(338) Problems concentrating in school**  
Options, 1: yes, very much, 2: sometimes, 5: no, never/rarely, 7: don't wish to answer, 8: don't know, 9: not known.
- 65      **(339) Special education.**  
Options, 1: yes, 5: no, 8: don't remember/know, 9: not known.
- 66      **(340) Bullied in school.**  
Options, 1: yes, often, 2: yes, rarely, 5: no, never, 8: don't remember/know, 9: not known.
- 67      **(341) Participation in follow-up study, if made**  
3 options, 1: yes, 2: no, 3: don't know.

The following last questions are rated by the interviewer

- 68      **(345) Danish language abilities**  
Options: 1: perfect (no accent), 2: very good ( a little accent), 3: Good (understandable with a strong accent), 4: poor, 5: does not speak Danish or only speak a little Danish, 8: don't know.
- 69      **(346) Has the subject answered questions 45-55 her-/himself.**  
Options; 1: yes, all of them, 2: yes, partly, 5: no, the interviewer has filled them in.